

Finite-dimensional vertex algebra modules over fixed point differential subfields

Kenichiro Tanabe*

Department of Mathematics
Hokkaido University
Kita 10, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810
Japan

ktanabe@math.sci.hokudai.ac.jp

Abstract

Let K be a differential field over $\mathbb{C}$ with derivation D , G a finite linear automorphism group over K which preserves D , and K^G the fixed point subfield of K under the action of G . We show that every finite-dimensional vertex algebra K^G -module is contained in some twisted vertex algebra K -module.

Keywords: vertex algebra; differential field

1 Introduction

In [2], Borcherds defined the notion of vertex algebras and showed that every commutative ring A with an arbitrary derivation D has a structure of vertex algebra. Every ring A -module naturally becomes a vertex algebra A -module. However, this does not imply that ring A -modules and vertex algebra A -modules are the same. In fact, a vertex algebra $\mathbb{Z}[z, z^{-1}]$ -module which is

*Partially supported by JSPS Grant-in-Aid for Scientific Research No. 20740002.

not a ring $\mathbb{Z}[z, z^{-1}]$ -module was given in [2, Section 8], where $\mathbb{Z}[z, z^{-1}]$ is the ring of Laurent polynomials over $\mathbb{Z}$. Moreover, in [9] for the polynomial ring $\mathbb{C}[s]$ in one variable s with derivation D , I obtained a necessary and sufficient condition on D that there exist finite-dimensional vertex algebra $\mathbb{C}[s]$ -modules which do not come from associative algebra $\mathbb{C}[s]$ -modules. Thus, in general vertex algebra A -modules and ring A -modules are certainly different.

Let K be a differential field with derivation D , G a finite linear automorphism group of K which preserves D , and K^G the fixed point subfield of K under the action of G . In this paper, we study vertex algebra K^G -modules. Here, let us recall the following conjecture on vertex operator algebras: let V be a vertex operator algebra and G a finite automorphism group of V . It is conjectured that under some conditions on V , every irreducible module over the fixed point vertex operator subalgebra V^G is contained in some irreducible g -twisted V -module for some $g \in G$ (cf.[4]). The motivation for studying vertex algebra K^G -modules is to investigate this conjecture for vertex algebras. In Theorem 1, I shall show that every finite-dimensional indecomposable vertex algebra K^G -module becomes a g -twisted vertex algebra K -module for some $g \in G$. Namely, the conjecture holds for all finite-dimensional vertex algebra K^G -modules in a stronger sense.

This paper is organized as follows. In Section 2 we recall some properties of vertex algebras and their modules. In Section 3 we show that every finite-dimensional indecomposable vertex algebra K^G -module becomes a g -twisted vertex algebra K -module for some $g \in G$. In Section 4 we give the classification of the finite-dimensional vertex algebra $\mathbb{C}(s)$ -modules where $\mathbb{C}(s)$ is the field of rational functions in one variable s . In Section 5 for all quadratic extensions K of $\mathbb{C}(s)$ and all finite-dimensional indecomposable vertex algebra $\mathbb{C}(s)$ -modules M obtained in Section 4, we study twisted vertex algebra K -module structures over M .

2 Preliminary

We assume that the reader is familiar with the basic knowledge on vertex algebras as presented in [2, 3, 7].

Throughout this paper, ζ_p is a primitive p -th root of unity for a positive integer p and $(V, Y, \mathbf{1})$ is a vertex algebra. Recall that V is the underlying vector space, $Y(\cdot, x)$ is the linear map from V to $(\text{End } V)[[x, x^{-1}]]$, and $\mathbf{1}$ is

the vacuum vector. Let $\mathcal{D}$ be the endomorphism of V defined by $\mathcal{D}v = v_{-2}\mathbf{1}$ for $v \in V$.

First, we recall some results in [2] for a vertex algebra constructed from a commutative associative algebra with a derivation.

Proposition 1. [2] *The following hold:*

- (1) *Let A be a commutative associative $\mathbb{C}$ -algebra with identity element $\mathbf{1}$ and D a derivation of A . For $a \in A$, define $Y(a, x) \in (\text{End } A)[[x]]$ by*

$$Y(a, x)b = \sum_{i=0}^{\infty} \frac{1}{i!} (D^i a) b x^i$$

for $b \in A$. Then, $(A, Y, \mathbf{1})$ is a vertex algebra.

- (2) *Let $(V, Y, \mathbf{1})$ be a vertex algebra such that $Y(u, x) \in (\text{End } V)[[x]]$ for all $u \in V$. Define a multiplication on V by $uv = u_{-1}v$ for $u, v \in V$. Then, V is a commutative associative $\mathbb{C}$ -algebra with identity element $\mathbf{1}$ and $\mathcal{D}$ is a derivation of V .*

Throughout the rest of this section, A is a commutative associative $\mathbb{C}$ -algebra with identity element $\mathbf{1}$ over $\mathbb{C}$ and D a derivation of A . Let $(A, Y, \mathbf{1})$ be the vertex algebra constructed from A and D in Proposition 1 and let (M, Y_M) be a module over vertex algebra A . We call M a *vertex algebra (A, D) -module* to distinguish between modules over vertex algebra A and modules over associative algebra A .

Proposition 2. [2] *The following hold:*

- (1) *Let M be an associative algebra A -module. For $a \in A$, define $Y_M(a, x) \in (\text{End}_{\mathbb{C}} M)[[x]]$ by*

$$Y(a, x)u = \sum_{i=0}^{\infty} \frac{1}{i!} (D^i a) u x^i$$

for $u \in M$. Then, (M, Y_M) is a vertex algebra (A, D) -module.

- (2) *Let (M, Y_M) be a vertex algebra (A, D) -module such that $Y(a, x) \in (\text{End}_{\mathbb{C}} M)[[x]]$ for all $a \in A$. Define an action of A on M by $au = a_{-1}u$ for $a \in A$ and $u \in M$. Then, M is an associative algebra A -module.*

Remark 1. (1) Let us consider the case of $D = 0$. Let (M, Y_M) be an arbitrary vertex algebra $(A, 0)$ -module. For all $a \in A$, since $0 = Y_M(0a, x) = dY_M(a, x)/dx$, we see that $Y_M(a, x)$ is constant. Thus, M is an associative algebra A -module by Proposition 2.

(2) Let us consider the case that A is finite-dimensional. Let (M, Y_M) be an arbitrary vertex algebra (A, D) -module. Suppose that there exists $a \in A$ and $u \in M$ such that $Y_M(a, x)u$ is not an element of $M[[x]]$. Since $Y_M(D^i a, x) = d^i Y_M(a, x)/dx^i$ for all $i \geq 0$, we see that $\{Y_M(D^i a, x)u \mid i = 0, 1, \dots\}$ is linearly independent. This contradicts that A is finite-dimensional. Thus, M is an associative algebra A -module by Proposition 2.

For a $\mathbb{C}$ -linear automorphism g of V of finite order p , set $V^r = \{u \in V \mid gu = \zeta_p^r u\}, 0 \leq r \leq p-1\}$. We recall the definition of g -twisted V -modules.

Definition 1. A g -twisted V -module M is a vector space equipped with a linear map

$$Y_M(\cdot, x): V \ni v \mapsto Y_M(v, x) = \sum_{i \in (1/p)\mathbb{Z}} v_i x^{-i-1} \in (\text{End}_{\mathbb{C}} M)[[x^{1/p}, x^{-1/p}]]$$

which satisfies the following four conditions:

- (1) $Y_M(u, x) = \sum_{i \in r/p + \mathbb{Z}} u_i x^{-i-1}$ for $u \in V^r$.
- (2) $Y_M(u, x)w \in M((x^{1/p}))$ for $u \in V$ and $w \in M$.
- (3) $Y_M(\mathbf{1}, x) = \text{id}_M$.
- (4) For $u \in V^r, v \in V^s, m \in r/T + \mathbb{Z}, n \in s/T + \mathbb{Z}$, and $l \in \mathbb{Z}$,

$$\begin{aligned} & \sum_{i=0}^{\infty} \binom{m}{i} (u_{l+i}v)_{m+n-i} \\ &= \sum_{i=0}^{\infty} \binom{l}{i} (-1)^i (u_{l+m-i}v_{n+i} + (-1)^{l+1} v_{l+n-i}u_{m+i}). \end{aligned}$$

The following result is well known (cf. [6, Proposition 4.8] and [9, Proposition 3]).

Proposition 3. *Let g be an automorphism of V of finite order p , M a vector space, and $Y_M(\cdot, x)$ a linear map from V to $(\text{End}_{\mathbb{C}} M)[[x^{1/p}, x^{-1/p}]]$ such that for all $0 \leq r \leq p-1$ and all $u \in V^r$, $Y_M(u, x) = \sum_{i \in r/p + \mathbb{Z}} u_i x^{-i-1}$. Then, (M, Y_M) is a g -twisted V -module if and only if the following five conditions hold:*

(M1) *For $u \in V$ and $w \in M$, $Y_M(u, x)w \in M((x^{1/p}))$.*

(M2) $Y_M(\mathbf{1}, x) = \text{id}_M$.

(M3) *For $u \in V^r, v \in V^s$, $m \in r/p + \mathbb{Z}$, and $n \in s/p + \mathbb{Z}$*

$$[u_m, v_n] = \sum_{i=0}^{\infty} \binom{m}{i} (u_i v)_{m+n-i}.$$

(M4) *For $u \in V^r, v \in V^s$, $m \in r/p + \mathbb{Z}$, and $n \in s/p + \mathbb{Z}$*

$$\sum_{i=0}^{\infty} \binom{m}{i} (u_{-1+i} v)_{m+n-i} = \sum_{i=0}^{\infty} (u_{-1+m-i} v_{n+i} + v_{-1+n-i} u_{m+i}).$$

(M5) *For $u \in V$, $Y_M(\mathcal{D}u, x) = dY_M(u, x)/dx$.*

Let $\mathcal{B}$ be a subset of A which generate A as a $\mathbb{C}$ -algebra and g a linear automorphism of A of finite order p . For a g -twisted A -module (M, Y_M) , we call (M, Y_M) a g -twisted vertex algebra (A, D) -module.

Lemma 4. *Let M be a vector space and $Y_M(\cdot, x)$ a linear map from A to $(\text{End}_{\mathbb{C}} M)[[x^{1/p}, x^{-1/p}]]$ such that for all $0 \leq r \leq p-1$ and all $a \in A^r$, $Y_M(a, x) = \sum_{i \in r/p + \mathbb{Z}} a_i x^{-i-1}$. Let $\mathcal{A}_M(A)$ denote the subalgebra of $\text{End}_{\mathbb{C}} M$ generated by all a_i where $a \in A$ and $i \in (1/p)\mathbb{Z}$. Suppose that M is a finitely generated $\mathcal{A}_M(A)$ -module.*

Then, (M, Y_M) is a g -twisted vertex algebra (A, D) -module if and only if the following five conditions hold:

(1) *For $a \in \mathcal{B}$, $Y_M(a, x) \in (\text{End}_{\mathbb{C}} M)((x^{1/p}))$.*

(2) $Y_M(\mathbf{1}, x) = \text{id}_M$.

(3) *For $a, b \in \mathcal{B}$ and $i, j \in (1/p)\mathbb{Z}$, $a_i b_j = b_j a_i$.*

(4) For $a \in \mathcal{B}$ and $b \in A$, $Y_M(ab, x) = Y_M(a, x)Y_M(b, x)$.

(5) For $a \in \mathcal{B}$, $Y_M(Da, x) = dY_M(a, x)/dx$.

In this case, $Y_M(\cdot, x)$ is a $\mathbb{C}$ -algebra homomorphism from A to $(\text{End}_{\mathbb{C}} M)((x^{1/p}))$.

Proof. We use Proposition 3. Note that for all $a, b \in A$ and all $i \geq 0$, we have $ab = a_{-1}b$ and $a_i b = 0$. Suppose that (M, Y_M) is a g -twisted vertex algebra (A, D) -module. We have (3) by (M3). Since M is a finitely generated $\mathcal{A}_M(A)$ -module, we have (1) by (M1) and (3). We have (4) by (M4), (1), and (3). The other conditions clearly hold.

Conversely, suppose that (M, Y_M) satisfies the conditions (1)–(5). It follows from (1), (2), and (4) that $Y_M(ab, x) = Y_M(a, x)Y_M(b, x)$ for all $a, b \in A$, namely $Y(\cdot, x)$ is a $\mathbb{C}$ -algebra homomorphism from A to $(\text{End}_{\mathbb{C}} M)((x^{1/p}))$. This shows (M3) by (3) and hence (M4). Let a, b be elements of A which satisfy (M5). Then,

$$\begin{aligned} Y_M(D(ab), x) &= Y_M((Da)b + a(Db), x) \\ &= Y_M(Da, x)Y_M(b, x) + Y_M(a, x)Y_M(Db, x) \\ &= \left(\frac{d}{dx}Y_M(a, x)\right)Y_M(b, x) + Y_M(a, x)\frac{d}{dx}Y_M(b, x) \\ &= \frac{d}{dx}(Y_M(a, x)Y_M(b, x)) \\ &= \frac{d}{dx}(Y_M(ab, x)). \end{aligned}$$

Since A is generated by $\mathcal{B}$, (M5) follows from (5). We conclude that (M, Y_M) is a g -twisted vertex algebra (A, D) -module. $\square$

For a g -twisted vertex algebra (A, D) -module (M, Y_M) and a linear automorphism h of A which preserves D , define $(M, Y_M) \circ h = (M \circ h, Y_{M \circ h})$ by $M \circ h = M$ as vector spaces and $Y_{M \circ h}(a, x) = Y_M(ha, x)$ for all $a \in A$. Then, $(M, Y_M) \circ h$ is an $h^{-1}gh$ -twisted vertex algebra (A, D) -module.

3 Finite-dimensional vertex algebra modules over fixed point differential subfields

Let K be a differential field over $\mathbb{C}$ with derivation D and let G be a finite $\mathbb{C}$ -linear automorphism group of K of order N which preserves D .

We fix a primitive element θ of K over K^G with the minimal polynomial $P(Z) = \sum_{i=0}^N P_i Z^i \in K^G[Z]$ in Z . For a finite-dimensional vertex algebra (K^G, D) -module (M, Y_M) , $g \in G$ of order p , and a linear map $\tilde{Y}(\cdot, x)$ from K to $(\text{End}_{\mathbb{C}} M)((x^{1/p}))$, we call $(M, \tilde{Y}_M)$ a *g -twisted vertex algebra* (K, D) -module structure over (M, Y_M) if $(M, \tilde{Y}_M)$ is a g -twisted vertex algebra (K, D) -module and if $\tilde{Y}(\cdot, x)|_{K^G} = Y(\cdot, x)$.

In this section, we shall show that every finite-dimensional indecomposable vertex algebra (K^G, D) -module (M, Y_M) has a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) for some $g \in G$. Here we give the outline of the proof. It follows from Lemma 4 that for any $g \in G$, to construct a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) is equivalent to construct a $\mathbb{C}$ -algebra homomorphism from K to $(\text{End}_{\mathbb{C}} M)((x^{1/|g|}))$ with some conditions. The basic idea to construct such a $\mathbb{C}$ -algebra homomorphism is to realize the Galois extension K/K^G in $(\text{End}_{\mathbb{C}} M)((x^{1/N}))$ where we identify K^G with its image under the homomorphism $Y_M(\cdot, x): K^G \rightarrow (\text{End}_{\mathbb{C}} M)((x))$. To do this, we first define a $\mathbb{C}$ -algebra homomorphism $\psi[K^G, (M, Y_M)]: K^G \rightarrow \mathbb{C}((x))$ for (M, Y_M) and denote by Q the image of $\psi[K^G, (M, Y_M)]$. It is well known that any finite extension of $\mathbb{C}((x))$ is $\mathbb{C}((x^{1/j}))$ for some positive integer j and $\Omega = \cup_{j=1}^{\infty} \mathbb{C}((x^{1/j}))$ is the algebraic closure of $\mathbb{C}((x))$ (cf. [5, Corollary 13.15]). These results enable us to construct the extension $\hat{K}$ of Q in Ω such that $\hat{K} \cong K$ as $\mathbb{C}$ -algebras. Using $\theta, P(Z)$, and $\hat{K}$, we can construct the desired extension corresponding to K in $(\text{End}_{\mathbb{C}} M)((x^{1/N}))$.

We introduce some notation. Let R be a commutative ring and let $\text{Mat}_n(R)$ denote the set of all $n \times n$ matrices with entries in R . Let E_n denote the $n \times n$ identity matrix and let E_{ij} denote the matrix whose (i, j) entry is 1 and all other entries are 0. Define $\Delta_k(R) = \{(x_{ij}) \in \text{Mat}_n(R) \mid x_{ij} = 0 \text{ if } i + k \neq j\}$ for $0 \leq k \leq n$. Then, for $a \in \Delta_k(R)$ and $b \in \Delta_l(R)$, we have $ab \in \Delta_{k+l}(R)$. For $X = (x_{ij}) \in \text{Mat}_n(R)$ and $k = 0, \dots, n-1$, define the matrix $X^{(k)} = \sum_{i=1}^n x_{i, i+k} E_{i, i+k} \in \Delta_k(R)$. For an upper triangular matrix X , we see that $X = \sum_{k=0}^{n-1} X^{(k)}$ and that the diagonal part of X is $X^{(0)}$. For a positive integer m and $H = \sum_{i \in (1/m)\mathbb{Z}} H_{(i)} x^i \in (\text{Mat}_n(R))[[x^{1/m}, x^{-1/m}]]$ with $H_{(i)} \in \text{Mat}_n(R)$, $H^{(k)}$ denotes $\sum_{i \in (1/m)\mathbb{Z}} H_{(i)}^{(k)} x^i \in \Delta_k(R)[[x^{1/m}, x^{-1/m}]]$ for $k = 0, \dots, n-1$. For an n -dimensional vector space M over $\mathbb{C}$, we sometimes identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ and $(\text{End}_{\mathbb{C}} M)[[x^{1/m}, x^{-1/m}]]$ with $(\text{Mat}_n(\mathbb{C}))[[x^{1/m}, x^{-1/m}]]$ by fixing a basis of M and use these symbols in the proofs of Theorem 1 and Theorem 2.

Let A be a commutative associative $\mathbb{C}$ -algebra, D a derivation of A , g a $\mathbb{C}$ -linear automorphism of A of finite order p . For a vector space W over $\mathbb{C}$ and a linear map $Y_W(\cdot, x)$ from A to $(\text{End } W)[[x^{1/p}, x^{-1/p}]]$, $\mathcal{A}_W(A)$ denotes the subalgebra of $\text{End } W$ generated by all coefficients of $Y_W(a, x)$ where a ranges over all elements of A . Let M be a finite-dimensional g -twisted vertex algebra (A, D) -module. Then, $\mathcal{A}_M(A)$ is a commutative $\mathbb{C}$ -algebra by Lemma 4 and M is a finite-dimensional $\mathcal{A}_M(A)$ -module. Let $\mathcal{J}_M(A)$ denote the Jacobson radical of $\mathcal{A}_M(A)$. Since $\mathcal{A}_M(A)$ is a finite-dimensional commutative $\mathbb{C}$ -algebra, the Wedderburn–Malcev theorem (cf.[8, Section 11.6]) says that $\mathcal{A}_M(A) = \oplus_{i=1}^m \mathbb{C}e_i \oplus \mathcal{J}_M(A)$ and hence

$$\mathcal{A}_M(A)((x^{1/p})) = \oplus_{i=1}^m \mathbb{C}((x^{1/p}))e_i \oplus \mathcal{J}_M(A)((x^{1/p}))$$

where $e_1, \dots, e_m$ are primitive orthogonal idempotents of $\mathcal{A}_M(A)$. In the case of $m = 1$, which is equivalent to $\mathcal{A}_M(A)$ being a local $\mathbb{C}$ -algebra, we shall often identify the subalgebra $\mathbb{C}((x^{1/p}))\text{id}$ of $\mathcal{A}_M(K)((x^{1/p}))$ with $\mathbb{C}((x^{1/p}))$. For $X \in \mathcal{A}_M(A)$, $X^{[0]}$ denotes the image of X under the projection

$$\begin{aligned} \mathcal{A}_M(A) &= \oplus_{i=1}^m \mathbb{C}e_i \oplus \mathcal{J}_M(A) \\ &\rightarrow \oplus_{i=1}^m \mathbb{C}e_i \cong \mathbb{C}^m \end{aligned} \tag{3.1}$$

and is called the *semisimple part* of X . For $H = \sum_{i \in (1/p)\mathbb{Z}} H_{(i)}x^i \in \mathcal{A}_M(A)((x^{1/p}))$ with $H_{(i)} \in \mathcal{A}_M(A)$, $H^{[0]}$ denotes $\sum_{i \in (1/p)\mathbb{Z}} H_{(i)}^{[0]}x^i \in \mathbb{C}((x^{1/p}))^{\oplus m}$ and is called the *semisimple part* of H . We denote by $\psi[A, (M, Y_M)]$ the $\mathbb{C}$ -algebra homomorphism $Y_M(\cdot, x)^{[0]}$ from A to $\mathbb{C}((x^{1/p}))^{\oplus m}$. Note that $\mathcal{J}_M(A)^n((x^{1/p})) = 0$ where $n = \dim_{\mathbb{C}} M$. If A is a field, then $\psi[A, (M, Y_M)]$ is an injective homomorphism from A to $\mathbb{C}((x^{1/p}))^{\oplus m}$. Since $\mathcal{A}_M(A)$ is commutative, we sometimes identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ by fixing a basis of M so that the representation matrix of each element of $\mathcal{A}_M(A)$ with respect to the basis is an upper triangular matrix. Under this identification, for $H \in \mathcal{A}_M(A)((x^{1/p})) \subset (\text{End}_{\mathbb{C}} M)((x^{1/p}))$ we see that $H^{[0]} = H^{(0)}$, which is the diagonal part of H defined above.

Let M be a finite-dimensional indecomposable vertex algebra (K^G, D) -module. Since $\mathcal{A}_M(K^G)$ is a commutative algebra, $\mathcal{A}_M(K^G)$ is a local algebra and hence $\mathcal{A}_M(K^G) = \mathbb{C}\text{id} \oplus \mathcal{J}_M(K^G)$. Let $(M, \tilde{Y}_M)$ be a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) . Since $\mathcal{A}_M(K^G)$ is a subalgebra of $\mathcal{A}_M(K)$, M is an indecomposable $\mathcal{A}_M(K)$ -module. Therefore, $\mathcal{A}_M(K)$ is local since $\mathcal{A}_M(K)$ is commutative. Thus, $\mathcal{A}_M(K) = \mathbb{C}\text{id} \oplus \mathcal{J}_M(K)$ and

hence

$$\psi[K, (M, \tilde{Y}_M)]|_{K^G} = \psi[K^G, (M, Y_M)]. \quad (3.2)$$

Symbol Q denotes the image of the homomorphism $\psi[K^G, (M, Y_M)] : K^G \rightarrow \mathbb{C}((x))$. For a polynomial $F(Z) \in K^G[Z]$, $\hat{F}(Z)$ denotes the image of $F(Z)$ under the map $Y_M(\cdot, x) : K^G[Z] \rightarrow (\mathcal{A}_M(K^G)((x)))[Z]$ and $\hat{F}^{[0]}(Z)$ denotes the image of $F(Z)$ under the map $\psi[K^G, (M, Y_M)] : K^G[Z] \rightarrow Q[Z] \subset \mathbb{C}((x))[Z]$. We write

$$\begin{aligned} \hat{F}(Z) &= \sum_{i \geq 0} \hat{F}_i(x) Z^i, \hat{F}_i(x) \in \mathcal{A}_M(K^G)((x)) \quad \text{and} \\ \hat{F}^{[0]}(Z) &= \sum_{i \geq 0} \hat{F}_i(x)^{[0]} Z^i, \hat{F}_i(x)^{[0]} \in Q. \end{aligned}$$

Since $P(Z)$ is an irreducible polynomial over K^G , so is $\hat{P}^{[0]}(Z)$ over Q .

Now we state our main theorem.

Theorem 1. *Let K be a differential field over $\mathbb{C}$ with derivation D , G a finite linear automorphism group of K which preserves D , and (M, Y_M) a non-zero finite-dimensional indecomposable vertex algebra (K^G, D) -module. Then, we have the following results:*

- (1) *M has a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) for some $g \in G$.*
- (2) *Let g be an element of G and let $(M, \tilde{Y}_M^1), (M, \tilde{Y}_M^2)$ be two g -twisted vertex algebra (K, D) -module structures over (M, Y_M) such that $\psi[K, (M, \tilde{Y}_M^1)] = \psi[K, (M, \tilde{Y}_M^2)]$. Then, $(M, \tilde{Y}_M^1) \cong (M, \tilde{Y}_M^2)$ as g -twisted vertex algebra (K, D) -modules.*
- (3) *Let g be an element of G and let $(M, \tilde{Y}_M)$ be a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) . Then, $\tilde{Y}_M \circ h, h \in G$, are all distinct homomorphisms from K to $(\text{End}_{\mathbb{C}} M)((x^{1/|g|}))$.*
- (4) *For $k = 1, 2$, let $g_k \in G$ of order p_k and let $(M, \tilde{Y}_M^k)$ be a g_k -twisted vertex algebra (K, D) -module structure over (M, Y_M) . Then, $(M, \tilde{Y}_M^1) \circ h \cong (M, \tilde{Y}_M^2)$ for some $h \in G$.*

Proof. Set $n = \dim_{\mathbb{C}} M$ and $|G| = N$. Let the notation be as above. Since K^G is a field, $\psi[K^G, (M, Y_M)]$ is an injective homomorphism from K^G to the field $\mathbb{C}((x))$ and hence $K^G \cong Q$ as $\mathbb{C}$ -algebras. It is well known that any finite extension of $\mathbb{C}((x))$ is $\mathbb{C}((x^{1/j}))$ for some positive integer j and $\Omega = \cup_{j=1}^{\infty} \mathbb{C}((x^{1/j}))$ is the algebraic closure of $\mathbb{C}((x))$ (cf. [5, Corollary 13.15]). The field $\mathbb{C}((x^{1/j}))$ becomes a Galois extension of $\mathbb{C}((x))$ whose Galois group is the cyclic group generated by the automorphism sending $x^{1/j}$ to $\zeta_j x^{1/j}$. Let $\hat{K}$ denote the splitting field for $\hat{P}^{[0]}(Z) \in Q[Z]$ in Ω and let $\mathcal{S}$ denote the set of all $\mathbb{C}$ -algebra isomorphisms from K to $\hat{K}$ whose restrictions to K^G are equal to $\psi[K^G, (M, Y_M)]$. Since θ is a primitive element of K over K^G , any map of $\mathcal{S}$ is uniquely determined by the image of θ under that map. Since K is the splitting field for $P(Z) \in K^G[Z]$, Galois theory implies that $|\mathcal{S}| = N$ and for any two $\phi_1, \phi_2 \in \mathcal{S}$ there exists $h \in G$ such that $\phi_1(ha) = \phi_2(a)$ for all $a \in K$.

Let g be an element of G and let $(M, \tilde{Y}_M)$ be a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) . By (3.2), for all root $a \in K$ of the polynomial $P(Z)$ we have

$$0 = \psi[K, (M, \tilde{Y}_M)](P(a)) = \hat{P}^{[0]}(\psi[K, (M, \tilde{Y}_M)](a)).$$

Since $\hat{K}$ is the splitting field for $\hat{P}^{[0]}(Z) \in Q[Z]$ in Ω , the image of K under the map $\psi[K, (M, \tilde{Y}_M)]$ is equal to $\hat{K}$ and hence $\psi[K, (M, \tilde{Y}_M)]$ is an element of $\mathcal{S}$.

We identify G with $\text{Gal}(\hat{K}/Q)$ throughout the argument below.

- (1) Let ϕ be an arbitrary element of $\mathcal{S}$. We use Lemma 4 by taking $\mathcal{B} = K^G \cup \{\theta\}$. We shall construct a $\mathbb{C}$ -algebra homomorphism $\tilde{Y}_M(\cdot, x)$ from $K \cong K^G[Z]/(P(Z))$ to $(\text{End}_{\mathbb{C}} M)((x^{1/N}))$. We first find $g \in G$ such that for all $b \in K$ with $gb = \zeta_p^i b, i \in \mathbb{Z}$, $\phi(b)$ is an element of $x^{-i/p} \mathbb{C}((x))$. Since K is a finite extension of K^G and Q is a subfield of $\mathbb{C}((x))$, Q is a subfield of $\hat{K} \cap \mathbb{C}((x))$ and $\hat{K} \mathbb{C}((x)) = \mathbb{C}((x^{1/p}))$ for some positive integer p . The isomorphism

$$\text{Gal}(\mathbb{C}((x^{1/p}))/\mathbb{C}((x))) \ni \sigma \mapsto \sigma|_{\hat{K}} \in \text{Gal}(\hat{K}/\hat{K} \cap \mathbb{C}((x))) \quad (3.3)$$

implies that $\text{Gal}(\hat{K}/\hat{K} \cap \mathbb{C}((x)))$ is the cyclic group of order p . Let $g \in G$ be a generator of $\text{Gal}(\hat{K}/\hat{K} \cap \mathbb{C}((x)))$ which is the homomorphic image of the element in $\text{Gal}(\mathbb{C}((x^{1/p}))/\mathbb{C}((x)))$ sending $x^{1/p}$ to $\zeta_p x^{1/p}$

under the isomorphism (3.3). We have the eigenspace decomposition $K = \bigoplus_{j=0}^{p-1} K^{(g;j)}$ for g where $K^{(g;j)} = \{a \in K \mid ga = \zeta_p^j a\}$. For $a \in K^{(g;i)}$ and $b \in K^{(g;j)}$, $1 \leq i, j \leq p-1$, taking two integers i_0, j_0 such that $i_0 i + j_0 j = (i, j)$, we have $g(a^{i_0} b^{j_0}) = \zeta_p^{(i,j)} a^{i_0} b^{j_0}$ where (i, j) is the greatest common divisor of i and j . This implies that we can take a nonzero element a of $K^{(g;1)}$ since the order of g is equal to p . We fix a nonzero element a of $K^{(g;1)}$. It follows from $\phi(a^p) \in \hat{K}^{(g)} = \hat{K} \cap \mathbb{C}((x))$ that $\phi(a)$ is a root of the polynomial $Z^p - \phi(a^p) \in \mathbb{C}((x))[Z]$. Thus, $\phi(a)$ is an element of $x^{-r/p} \mathbb{C}((x))$ for some integer r . For all $i = 1, \dots, p-1$, it follows by $a^i \notin K^{(g)}$ that $\phi(a^i) \notin \hat{K}^{(g)} = \hat{K} \cap \mathbb{C}((x))$. Therefore, we have $(r, p) = 1$. Taking two integers γ, δ such that $\gamma r + \delta p = 1$ and replacing a by a^γ , we have $ga = \zeta_p^\gamma a$ and $\phi(a) \in x^{-1/p} \mathbb{C}((x))$. Since $(\gamma, p) = 1$, by replacing g by a suitable power of g , we have $ga = \zeta_p a$ and $\phi(a) \in x^{-1/p} \mathbb{C}((x))$. For all $b \in K$ with $gb = \zeta_p^i b$, since $a^{-i} b$ is an element of $K^{(g)} = \hat{K} \cap \mathbb{C}((x))$, we have $\phi(b)$ is an element of $x^{-i/p} \mathbb{C}((x))$. Set $T(x)^{[0]} = \phi(\theta) \in \hat{K}$, which is a root of $\hat{P}^{[0]}(Z) \in Q[Z]$. Since $\hat{P}^{[0]}(Z)$ is the image of $P(Z)$ under the map $\psi[K^G, (M, Y_M)]$, it is a separable polynomial and hence $(d\hat{P}^{[0]}/dZ)(T(x)^{[0]})$ is not zero. We set

$$\begin{aligned} \hat{P}_i(x)^{[1]} &= \hat{P}_i(x) - \hat{P}_i(x)^{[0]} \in \mathcal{J}_M(K^G)((x)) \quad \text{and} \\ \hat{P}_i(x)^{[k]} &= 0 \in \mathcal{J}_M(K^G)^k((x)) \end{aligned}$$

for all $i = 0, \dots, N$ and $k = 2, 3, \dots$ for convenience. For $k = 1, 2, \dots, n-1$ we inductively define $T(x)^{[k]} \in \mathcal{J}_M(K^G)^k((x^{1/p}))$ by

$$\begin{aligned} T(x)^{[k]} &= -\left(\frac{d\hat{P}^{[0]}}{dZ}(T(x)^{[0]})\right)^{-1} \\ &\quad \times \sum_{i=0}^N \sum_{j_0=0}^k \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{[j_0]} T(x)^{[j_1]} \dots T(x)^{[j_i]}. \end{aligned} \quad (3.4)$$

Set $T(x) = \sum_{k=0}^{n-1} T(x)^{[k]} \in \mathcal{A}_M(K^G)((x^{1/p}))$. It follows from the definition of $T(x)$ that $T(x)^{[0]}$ is exactly the semisimple part of $T(x)$ defined

arter (3.1). Since $\mathcal{J}_M(K^G)^n((x)) = 0$, we have

$$\begin{aligned}
\hat{P}(T(x)) &= \sum_{i=0}^N \hat{P}_i(x)(T(x))^i \\
&= \sum_{i=0}^N \left(\sum_{j=0}^{n-1} \hat{P}_i(x)^{[j]} \right) \left(\sum_{k=0}^{n-1} T(x)^{[k]} \right)^i \\
&= \sum_{k=0}^{n-1} \sum_{i=0}^N \sum_{\substack{0 \leq j_0, j_1, \dots, j_i \leq n \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{[j_0]} T(x)^{[j_1]} \dots T(x)^{[j_i]} \\
&= \hat{P}^{[0]}(T(x)^{[0]}) \\
&\quad + \sum_{k=1}^{n-1} \sum_{i=0}^N \sum_{\substack{0 \leq j_0, j_1, \dots, j_i \leq k \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{[j_0]} T(x)^{[j_1]} \dots T(x)^{[j_i]} \\
&= 0 + \sum_{k=1}^{n-1} \left(T(x)^{[k]} \frac{d\hat{P}^{[0]}}{dZ}(T(x)^{[0]}) \right) \\
&\quad + \sum_{i=0}^N \sum_{j_0=0}^k \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{[j_0]} T(x)^{[j_1]} \dots T(x)^{[j_i]} \\
&= 0.
\end{aligned} \tag{3.5}$$

Thus, we have an injective homomorphism $\tilde{Y}_M(\cdot, x)$ from $K \cong K^G[Z]/(P(Z))$ to $\mathcal{A}_M(K^G)((x^{1/p}))$ sending θ to $T(x)$. Since $T(x)$ is an element of $\mathcal{A}_M(K^G)((x^{1/p}))$, the $\mathbb{C}$ -algebra $\mathcal{A}_M(K)$ for $\tilde{Y}_M(\cdot, x)$ is equal to $\mathcal{A}_M(K^G)$. In particular, $\mathcal{A}_M(K)$ is a commutative $\mathbb{C}$ -algebra. Moreover, it follows from $\phi(\theta) = T(x)^{[0]}$ that $\psi[K, (\tilde{Y}_M, M)] = \phi$.

Let b be an element of K with the minimal polynomial $R(Z) \in K^G[Z]$ over K^G and let $B(x)$ denotes $\tilde{Y}_M(b, x)$. We write $R(Z) = \sum_{i=0}^m R_i Z^i \in K^G[Z]$, $R_i \in K^G$. We shall show that if $B(x)^{[0]}$ is an element of $\mathbb{C}((x)) \text{id}$, then $B(x)$ is an element of $(\text{End}_{\mathbb{C}} M)((x))$. By definition, $\hat{R}(B(x)) = 0$ in $(\text{End}_{\mathbb{C}} M)((x^{1/p}))$ and $\hat{R}^{[0]}(B(x)^{[0]}) = 0$ in Ωid . We identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ by fixing a basis of M so that the representation matrix of each element of $\mathcal{A}_M(K^G)$ with respect to the basis is an upper triangular matrix. We use the expansion $B(x) = \sum_{k=0}^{n-1} B(x)^{(k)}$, $B(x)^{(k)} \in \Delta_k(\mathbb{C}((x^{1/p})))$. We recall $B(x)^{[0]} = B(x)^{(0)}$

and $\hat{R}_i(x)^{[0]} = \hat{R}_i(x)^{(0)}$, $i = 0, \dots, m$. It may be possible that $B(x)^{(k)}$ and $\hat{R}_i(x)^{(k)}$ are not elements of $\mathcal{A}_M(K^G)((x^{1/p}))$ for $k = 1, \dots, n-1$. Since $B(x)^{(0)}$ and $\hat{R}_i(x)^{(0)}$ are elements of $\mathbb{C}((x^{1/p}))E_n$, they commute any element of $(\text{End}_{\mathbb{C}} M)((x^{1/p}))$. We denote the diagonal part of $\hat{R}(Z)$ by $\hat{R}^{(0)}(Z)$, namely, $\hat{R}^{(0)}(Z) = \sum_{i=0}^m \hat{R}_i(x)^{(0)} Z^i$. Under the identification of $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ above, $\hat{R}^{(0)}(Z)$ is equal to $\hat{R}^{[0]}(Z)$. The same computation as (3.5) shows

$$\begin{aligned}
0 &= \hat{R}(B(x)) = \sum_{i=0}^m \hat{R}_i(x) (B(x))^i \\
&= \sum_{i=0}^m \left(\sum_{j=0}^{n-1} \hat{R}_i(x)^{(j)} \right) \left(\sum_{k=0}^{n-1} B(x)^{(k)} \right)^i \\
&= \sum_{k=0}^{n-1} \sum_{i=0}^m \sum_{\substack{0 \leq j_0, j_1, \dots, j_i \leq n \\ j_0 + j_1 + \dots + j_i = k}} \hat{R}_i(x)^{(j_0)} B(x)^{(j_1)} \dots B(x)^{(j_i)} \\
&= \hat{R}^{(0)}(B(x)^{(0)}) \\
&\quad + \sum_{k=1}^{n-1} \sum_{i=0}^m \sum_{\substack{0 \leq j_0, j_1, \dots, j_i \leq k \\ j_0 + j_1 + \dots + j_i = k}} \hat{R}_i(x)^{(j_0)} B(x)^{(j_1)} \dots B(x)^{(j_i)} \\
&= \sum_{k=1}^{n-1} \left(B(x)^{(k)} \frac{d\hat{R}^{(0)}}{dZ} (B(x)^{(0)}) \right. \\
&\quad \left. + \sum_{i=0}^m \sum_{j_0=0}^k \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \hat{R}_i(x)^{(j_0)} B(x)^{(j_1)} \dots B(x)^{(j_i)} \right). \quad (3.6)
\end{aligned}$$

Since $\hat{R}^{(0)}(Z) \in Q[Z]$ is the image of $R(Z)$ under the map $\psi[K^G, (M, Y_M)]$, $\hat{R}^{(0)}(Z)$ is a separable polynomial and hence $(d\hat{R}^{(0)}/dZ)(B(x)^{(0)})$ is not zero. Thus, we have

$$\begin{aligned}
B(x)^{(k)} &= - \left(\frac{d\hat{R}^{(0)}}{dZ} (B(x)^{(0)}) \right)^{-1} \\
&\quad \times \sum_{i=0}^m \sum_{j_0=0}^k \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \hat{R}_i(x)^{(j_0)} B(x)^{(j_1)} \dots B(x)^{(j_i)}.
\end{aligned}$$

Since $B(x)^{(0)} = \phi(b) \in \mathbb{C}((x))E_n$ and $\hat{R}_i(x) \in (\text{End}_{\mathbb{C}} M)((x))$, it follows by induction on k that $B(x)^{(k)}$ is an element of $(\text{End}_{\mathbb{C}} M)((x))$. We conclude that $B(x)$ is an element of $(\text{End}_{\mathbb{C}} M)((x))$.

For all $b \in K$ with $gb = \zeta_p^i b$, we shall show that $\tilde{Y}_M(b, x)$ is an element of $x^{-i/p}(\text{End}_{\mathbb{C}} M)((x))$. Set $B(x) = \tilde{Y}_M(b, x)$ and $C(x) = B(x)^p$. It follows from $b^p \in K^{(g)}$ that $C(x)^{[0]} = (B(x)^{[0]})^p$ is an element of $\hat{K}^{(g)} = \hat{K} \cap \mathbb{C}((x))$. The argument above shows $C(x)$ is an element of $(\text{End}_{\mathbb{C}} M)((x))$. We identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ by fixing a basis of M so that the representation matrix of each element of $\mathcal{A}_M(K^G)$ with respect to the basis is an upper triangular matrix. We use the expansion $B(x) = \sum_{k=0}^{n-1} B(x)^{(k)}$, $B(x)^{(k)} \in \Delta_k(\mathbb{C}((x^{1/p})))$. We have already seen in the first part of the proof of (1) that $B(x)^{(0)} = \phi(b)$ is an element of $x^{-i/p}(\text{End}_{\mathbb{C}} M)((x))$. By $C(x) = B(x)^p$, the same computation as (3.6) shows

$$\begin{aligned} B(x)^{(k)} &= -p^{-1}(B(x)^{(0)})^{-p+1} \\ &\quad \times (C(x)^{(k)} + \sum_{\substack{0 \leq j_1, \dots, j_p < k \\ j_1 + \dots + j_p = k}} B(x)^{(j_1)} \dots B(x)^{(j_p)}). \end{aligned}$$

for all $k = 1, \dots, n-1$. It follows by induction on k that $B(x)^{(k)}$ is an element of $x^{-i/p}(\text{End}_{\mathbb{C}} M)((x))$. We conclude that $B(x)$ is an element of $x^{-i/p}(\text{End}_{\mathbb{C}} M)((x))$.

We shall show $\tilde{Y}_M(D\theta, x) = d\tilde{Y}_M(\theta, x)/dx$. It follows from $P(\theta) = 0$ that

$$0 = D(P(\theta)) = \sum_{i=0}^N (DP_i)\theta^i + \left(\frac{dP}{dZ}(\theta)\right)(D\theta).$$

and hence

$$0 = \sum_{i=0}^N \tilde{Y}_M(DP_i, x) \tilde{Y}_M(\theta, x)^i + \tilde{Y}_M\left(\frac{dP}{dZ}(\theta), x\right) \tilde{Y}_M(D\theta, x). \quad (3.7)$$

We also have

$$\begin{aligned}
0 &= \frac{d}{dx} \tilde{Y}_M(P(\theta), x) \\
&= \frac{d}{dx} \sum_{i=0}^N \tilde{Y}_M(P_i, x) \tilde{Y}_M(\theta, x)^i \\
&= \sum_{i=0}^N \frac{d\tilde{Y}_M(P_i, x)}{dx} \tilde{Y}_M(\theta, x)^i + \sum_{i=0}^N \tilde{Y}_M(P_i, x) i \tilde{Y}_M(\theta, x)^{i-1} \frac{\tilde{Y}_M(\theta, x)}{dx} \\
&= \sum_{i=0}^N \frac{dY_M(P_i, x)}{dx} \tilde{Y}_M(\theta, x)^i + \tilde{Y}_M\left(\frac{dP}{dZ}(\theta), x\right) \tilde{Y}_M \frac{\tilde{Y}_M(\theta, x)}{dx}. \tag{3.8}
\end{aligned}$$

Since $P(Z) \in K^G[Z]$ is a separable polynomial in Z , $(dP/dZ)(\theta)$ is not zero and hence $\tilde{Y}_M((dP/dZ)(\theta), x)$ is an invertible element of $(\text{End}_{\mathbb{C}} M)((x^{1/p}))$. Since $Y_M(DP_i, x) = dY_M(P_i, x)/dx$ for all i , we have $\tilde{Y}_M(D\theta, x) = d\tilde{Y}_M(\theta, x)/dx$ by (3.7) and (3.8).

By Lemma 4, we conclude that $(M, \tilde{Y}_M)$ is a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) .

- (2) We denote the order of g by p and $\psi[K, (M, \tilde{Y}_M^1)] = \psi[K, (M, \tilde{Y}_M^2)]$ by Ψ for simplicity. We recall Ψ is an element of $\mathcal{S}$ as mentioned before starting the proof of (1).

Let $(M, \tilde{Y}_M)$ be a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) constructed in (1) by taking Ψ as ϕ . It is enough to show that $\tilde{Y}_M = \tilde{Y}_M^1$. We denote $\mathcal{A}_M(K)$ for $(M, \tilde{Y}_M)$ by $\mathcal{A}^0$ and $\mathcal{A}_M(K)$ for $(M, \tilde{Y}_M^1)$ by $\mathcal{A}^1$. We have seen in (1) that $\mathcal{A}^0 = \mathcal{A}_M(K^G)$. Thus, $\mathcal{A}^0$ is a subalgebra of $\mathcal{A}^1$. We identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ by fixing a basis of M so that the representation matrix of each element of $\mathcal{A}^1$ with respect to the basis is an upper triangular matrix. We denote $\tilde{Y}_M^1(\theta, x)$ by $U(x)$. We use the expansions

$$\begin{aligned}
U(x) &= \sum_{k=0}^{n-1} U(x)^{(k)}, \quad U(x)^{(k)} \in \Delta_k(\mathbb{C}((x^{1/p}))) \text{ and} \\
\hat{P}_i(x) &= \sum_{k=0}^{n-1} \hat{P}_i(x)^{(k)}, \quad \hat{P}_i(x)^{(k)} \in \Delta_k(\mathbb{C}((x^{1/p}))).
\end{aligned}$$

We denote the diagonal part of $\hat{P}(Z)$ by $\hat{P}^{(0)}(Z)$, namely, $\hat{P}^{(0)}(Z) = \sum_{i=0}^N \hat{P}_i(x)^{(0)} Z^i$. Under the identification of $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ above, $\hat{P}^{(0)}(Z)$ is equal to $\hat{P}^{[0]}(Z)$. Note that we do not assume $U(x) \in \mathcal{A}_M(K^G)((x^{1/p}))$. We recall $\Psi(\theta) = U(x)^{(0)}$ by definition. We have $\hat{P}^{(0)}(U(x)^{(0)}) = \Psi(P(\theta)) = 0$ and $(d\hat{P}^{(0)}/dZ)(U(x)^{(0)})$ is not zero since $\hat{P}^{[0]}(Z)$ is a separable polynomial. The same computation as (3.6) shows

$$\begin{aligned}
0 &= \hat{P}(U(x)) \\
&= \hat{P}^{(0)}(U(x)^{(0)}) \\
&\quad + \sum_{k=1}^{n-1} \sum_{i=0}^N \sum_{\substack{0 \leq j_0, j_1, \dots, j_i \leq k \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{(j_0)} U(x)^{(j_1)} \dots U(x)^{(j_i)} \\
&= \sum_{k=1}^{n-1} \left(U(x)^{(k)} \frac{d\hat{P}^{(0)}}{dZ}(U(x)^{(0)}) \right. \\
&\quad \left. + \sum_{i=0}^N \sum_{j_0=0}^k \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{(j_0)} U(x)^{(j_1)} \dots U(x)^{(j_i)} \right)
\end{aligned}$$

and hence

$$\begin{aligned}
U(x)^{(k)} &= - \left(\frac{d\hat{P}^{[0]}}{dZ}(\Psi(\theta)) \right)^{-1} \\
&\quad \times \sum_{i=0}^N \sum_{j_0=0}^k \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \hat{P}_i(x)^{(j_0)} U(x)^{(j_1)} \dots U(x)^{(j_i)}.
\end{aligned}$$

for all $k = 1, \dots, n-1$. It follows by induction on k that $U(x) = \sum_{k=0}^{n-1} U(x)^{(k)}$ is uniquely determined by $\Psi(\theta)$ and $\hat{P}(Z)$. By definition of Ψ , we have $\tilde{Y}_M(\theta, x) = U(x) = \tilde{Y}_M^1(\theta, x)$ and hence $\tilde{Y}_M = \tilde{Y}_M^1$.

- (3) Let $h \in G$ with $h \neq 1$. Since $h^{-1}(\theta) \neq \theta$ and $\tilde{Y}_{M \circ h}(h^{-1}\theta, x) = \tilde{Y}_M(\theta, x)$, $\tilde{Y}_{M \circ h}(\cdot, x)$ is distinct from $\tilde{Y}_M(\cdot, x)$. This implies that $\tilde{Y}_M \circ h, h \in G$, are all distinct homomorphisms from K to $(\text{End}_{\mathbb{C}} M)((x^{1/|g|}))$.
- (4) For $k = 1, 2$, let $g_k \in G$ of order p_k and let $(M, \tilde{Y}_M^k)$ be a g_k -twisted vertex algebra (K, D) -module structure over (M, Y_M) . We denote $\psi[K, (M, \tilde{Y}_M^k)]$ by ψ_k and $\psi[K^G, (M, Y_M)]$ by ψ briefly.

Since $\psi_1, \psi_2 \in \mathcal{S}$, there exists $h \in G$ such that $\psi_1(ha) = \psi_2(a)$ for all $a \in K$. By definition of $(M, \tilde{Y}_M^1) \circ h$, we have $\psi[K, (M, \tilde{Y}_M^1) \circ h](a) = \psi_1(ha) = \psi_2(a)$ for all $a \in K$. We conclude that $(M, \tilde{Y}_M^1) \circ h \cong (M, \tilde{Y}_M^2)$ by (2).

□

4 Finite-dimensional vertex algebra $\mathbb{C}(s)$ -modules

Throughout the rest of this paper, $\mathbb{C}(s)$ is the field of rational functions in one variable s . In this section we classify the finite-dimensional vertex algebra $\mathbb{C}(s)$ -modules. We use the notation introduced in Section 3. It is easy to see that every non-zero derivation D of $\mathbb{C}(s)$ can be expressed as $D = (p(s)/q(s))d/ds$, where $p(s)$ and $q(s)$ are non-zero coprime elements of $\mathbb{C}[s]$. We write

$$p(s) = \sum_{i=L_p}^{N_p} p_i s^i \quad \text{and} \quad q(s) = \sum_{i=L_q}^{N_q} q_i s^i$$

where $p_{L_p}, p_{N_p}, q_{L_q}, q_{N_q}$ are all non-zero complex numbers.

The following lemma is a corollary of Lemma 4.

Lemma 5. *Let the notation be as above. Let M be a finite-dimensional vector space and let $S(x) = \sum_{i \in \mathbb{Z}} S_{(i)} x^i$ be an element of $(\text{End}_{\mathbb{C}} M)((x))$. Then, there exists a vertex algebra $(\mathbb{C}(s), D)$ -module (M, Y_M) with $Y_M(s, x) = S(x)$ if and only if the following three conditions hold:*

- (i) *For all $\alpha \in \mathbb{C}$, $S(x) - \alpha$ is an invertible element of $(\text{End}_{\mathbb{C}} M)((x))$.*
- (ii) *For all $i, j \in \mathbb{Z}$, $S_{(i)} S_{(j)} = S_{(j)} S_{(i)}$.*
- (iii) *$dS(x)/dx = p(S(x))/q(S(x))$.*

In this case, for $u(s) \in \mathbb{C}(s)$ we have $Y_M(u(s), x) = u(S(x))$ and hence (M, Y_M) is uniquely determined by $S(x)$.

Proof. We use Lemma 4 by taking $\mathcal{B} = \{s\} \cup \{(s - \alpha)^{-1} \mid \alpha \in \mathbb{C}\}$. Suppose that (M, Y_M) is a vertex algebra $(\mathbb{C}(s), D)$ -module. Then, $Y_M(\cdot, x)$ is a $\mathbb{C}$ -algebra homomorphism from $\mathbb{C}(s)$ to $(\text{End}_{\mathbb{C}} M)((x))$. For all $\alpha \in \mathbb{C}$, since

$s - \alpha$ is an invertible element in $\mathbb{C}(s)$, so is $Y_M(s - \alpha, x) = S(x) - \alpha$ in $(\text{End}_{\mathbb{C}} M)((x))$. The other conditions in (ii) and (iii) are clearly hold.

Conversely, suppose that (M, Y_M) satisfies the conditions (i)–(iii). Since $S(x) - \alpha$ is an invertible element of $(\text{End}_{\mathbb{C}} M)((x))$ for all $\alpha \in \mathbb{C}$, we obtain $q(S(x)) \neq 0$ and we can define the $\mathbb{C}$ -algebra homomorphism $Y_M: \mathbb{C}(s) \ni u(s) \mapsto u(S(x)) \in (\text{End}_{\mathbb{C}} M)((x))$. Since for all $\alpha \in \mathbb{C}$ each coefficient of $x^j, j \in \mathbb{Z}$ in the expansion of $(S(x) - \alpha)^{-1}$ is a polynomial in $\{S_{(k)} \mid k \in \mathbb{Z}\}$, $\mathcal{A}_M(\mathbb{C}(s))$ is commutative. For all $\alpha \in \mathbb{C}$ we have

$$\begin{aligned} Y_M(D((s - \alpha)^{-1}), x) &= Y_M(-(Ds)(s - \alpha)^{-2}, x) \\ &= -Y_M(Ds, x)(Y_M(s, x) - \alpha)^{-2} \\ &= \frac{d}{dx}(Y_M(s, x) - \alpha)^{-1}. \end{aligned}$$

We conclude that (M, Y_M) is a vertex algebra $(\mathbb{C}(s), D)$ -module. $\square$

Note that there is no nontrivial finite-dimensional associative algebra $\mathbb{C}(s)$ -module since $\mathbb{C}(s)$ is an infinite-dimensional $\mathbb{C}$ -vector space and $\mathbb{C}(s)$ is a field.

Let M be a finite-dimensional vector space over $\mathbb{C}$. For $X \in \text{End}_{\mathbb{C}} M$, $X^{[0]}$ denotes the semisimple part of X and $X^{[1]}$ denotes the nilpotent part of X . For $H(x) = \sum_{i \in \mathbb{Z}} H_{(i)} x^i \in (\text{End}_{\mathbb{C}} M)[[x, x^{-1}]]$, $H(x)^{[0]}$ denotes $\sum_{i \in \mathbb{Z}} H_{(i)}^{[0]} x^i$ and $H(x)^{[1]}$ denotes $\sum_{i \in \mathbb{Z}} H_{(i)}^{[1]} x^i$. We call $H(x)^{[0]}$ the semisimple part of $H(x)$. For $H(x) = \sum_{i=L}^{\infty} H_{(i)} x^i \in (\text{End}_{\mathbb{C}} M)((x))$ with $H_{(L)} \neq 0$, $\text{ld}(H(x))$ denotes L and $\text{lc}(H(x))$ denotes $H_{(L)}$.

For a finite-dimensional indecomposable vertex algebra $\mathbb{C}(s)$ -module (M, Y_M) , we denote $Y_M(s, x)$ by $S(x)$. Let J_n denote the following $n \times n$ matrix:

$$J_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ \vdots & & & \ddots & 1 \\ 0 & \cdots & \cdots & \cdots & 0 \end{pmatrix}.$$

Theorem 2. *Let the notation be as above. Let α be a non-zero complex number. We have the following results:*

- (1) *There exists a non-zero finite-dimensional indecomposable vertex algebra $(\mathbb{C}(s), D)$ -module M with $\text{ld}(S(x)^{[0]}) > 0$ and with $\text{lc}(S(x)^{[0]}) = \alpha$*

if and only if $p(0)q(0) \neq 0$ and $\alpha = p(0)/q(0)$. Moreover, in this case $\text{ld}(S(x)^{[0]}) = 1$ and $S(x) \in (\text{End}_{\mathbb{C}} M)[[x]]$.

- (2) There exists a non-zero finite-dimensional indecomposable vertex algebra $(\mathbb{C}(s), D)$ -module M with $\text{ld}(S(x)^{[0]}) = 0$ and with $\text{lc}(S(x)^{[0]}) = \alpha$ if and only if $p(\alpha)q(\alpha) \neq 0$. Moreover, in this case $S(x) \in (\text{End}_{\mathbb{C}} M)[[x]]$ and $S_{(1)}^{[0]} = p(\alpha)/q(\alpha)$.
- (3) There exists a non-zero finite-dimensional indecomposable vertex algebra $(\mathbb{C}(s), D)$ -module M with $\text{ld}(S(x)^{[0]}) < 0$ and with $\text{lc}(S(x)^{[0]}) = \alpha$ if and only if $\deg p = \deg q + 2$ and $\alpha = -q_{N_q}/p_{N_p}$. Moreover, in this case $\text{ld}(S(x)^{[0]}) = -1$.

In each case, for all positive integers n , there exists a unique n -dimensional indecomposable vertex algebra $(\mathbb{C}(s), D)$ -module which satisfies the conditions up to isomorphism.

Proof. (1) Let n be a positive integer and let (M, Y_M) be an n -dimensional indecomposable vertex algebra $(\mathbb{C}(s), D)$ -module and let $S(x)$ denotes $Y_M(s, x)$. Since M is indecomposable, $S(x)^{[0]}$ is an element of $\mathbb{C}((x)) \text{ id}$. If $S(x)^{[0]} = 0$, then $S(x)$ is nilpotent. This is impossible since $\mathbb{C}(s)$ is a field. Thus, we write $S(x)^{[0]} = \sum_{i=L}^{\infty} S_{(i)}^{[0]} x^i$ where $L = \text{ld}(S(x)^{[0]})$. Since $D = (p(s)/q(s))d/ds$, we have

$$\frac{dS(x)}{dx} = q(S(x))^{-1}p(S(x)) \quad (4.1)$$

or equivalently

$$q(S(x)) \frac{dS(x)}{dx} = p(S(x)). \quad (4.2)$$

Since $q(S(x))$ is an invertible element of $(\text{End}_{\mathbb{C}} M)((x))$, $q(S(x)^{[0]})$ is not zero. Taking the semisimple part of (4.2), we have

$$q(S(x)^{[0]}) \frac{dS(x)^{[0]}}{dx} = p(S(x)^{[0]}). \quad (4.3)$$

Suppose that L is a positive integer, namely $S(x)^{[0]}$ is an element of $x\mathbb{C}[[x]]$. Then, $S_{(0)}^{[0]} = 0$ and hence $S_{(0)}$ is a nilpotent element of $\text{End}_{\mathbb{C}} M$.

We shall show $p(0)q(0) \neq 0$, $\text{lc}(S(x)^{[0]}) = 1$, and $\text{lc}(S(x)^{[0]}) = p(0)/q(0)$.

In (4.3), the term with the lowest degree of the left-hand side is $q_{L_q} L(S_{(L)}^{[0]})^{L_q+1} x^{L(L_q+1)-1}$

and the term with the lowest degree of the right-hand side is $p_{L_p} (S_{(L)}^{[0]})^{L_p} x^{LL_p}$.

Comparing these terms, we have $L(L_p - L_q - 1) = -1$. Therefore, $L = 1$ and $L_p = L_q$. We have $L_p = L_q = 0$ since $p(x)$ and $q(x)$ are coprime. Thus, both p_0, q_0 are not zero. Comparing the coefficients of these terms, we also have $S_{(1)}^{[0]} = p_0/q_0 = p(0)/q(0)$.

We shall show that $S(x)$ is an element of $(\text{End}_{\mathbb{C}} M)[[x]]$. In order to do that, we identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ by fixing a basis of M so that the representation matrix of each element of $\mathcal{A}_M(\mathbb{C}(s))$ with respect to the basis is upper triangular matrix for a while. We write the expansion $S(x) = \sum_{k=0}^{n-1} S(x)^{(k)}$, $S(x)^{(k)} \in \Delta_k(\mathbb{C}((x)))$ as in Section 3. We recall that $S(x)^{[0]}$ is equal to the diagonal part $S(x)^{(0)}$. We need to expand $(dS(x)/dx)q(S(x))$ and $p(S(x))$. By the same computation as (3.5), we have

$$\begin{aligned}
& \frac{dS(x)}{dx} q(S(x)) \\
&= \sum_{j_0=0}^{n-1} \frac{dS(x)^{(j_0)}}{dx} \left(\sum_{i=0}^{N_q} q_i \sum_{0 \leq j_1, \dots, j_i \leq n-1} S(x)^{(j_1)} \dots S(x)^{(j_i)} \right) \\
&= \sum_{i=0}^{N_q} q_i \sum_{0 \leq j_0, j_1, \dots, j_i \leq n-1} \frac{dS(x)^{(j_0)}}{dx} S(x)^{(j_1)} \dots S(x)^{(j_i)} \\
&= \sum_{k=0}^{n-1} \sum_{i=0}^{N_q} q_i \sum_{\substack{0 \leq j_0, j_1, \dots, j_i \leq k \\ j_0 + j_1 + \dots + j_i = k}} \frac{dS(x)^{(j_0)}}{dx} S(x)^{(j_1)} \dots S(x)^{(j_i)} \\
&= \frac{dS(x)^{(0)}}{dx} q(S(x)^{(0)}) \\
&\quad + \sum_{k=1}^{n-1} \left(\frac{dS(x)^{(k)}}{dx} q(S(x)^{(0)}) + \frac{dS(x)^{(0)}}{dx} \frac{dq}{ds}(S(x)^{(0)}) S(x)^{(k)} \right) \\
&\quad + \sum_{i=0}^{N_q} q_i \sum_{\substack{0 \leq j_0, j_1, \dots, j_i < k \\ j_0 + j_1 + \dots + j_i = k}} \frac{dS(x)^{(j_0)}}{dx} S(x)^{(j_1)} \dots S(x)^{(j_i)}
\end{aligned}$$

and

$$\begin{aligned}
p(S(x)) &= p(S(x)^{(0)}) + \sum_{k=1}^{n-1} \left(\frac{dp}{ds}(S(x)^{(0)}) S(x)^{(k)} \right. \\
&\quad \left. + \sum_{i=0}^{N_p} p_i \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_1 + \dots + j_i = k}} S(x)^{(j_1)} \dots S(x)^{(j_i)} \right).
\end{aligned}$$

Thus, it follows from (4.2) that for all $k = 1, 2, \dots, n-1$

$$\begin{aligned}
&\frac{dS(x)^{(k)}}{dx} \\
&= q(S(x)^{(0)})^{-1} \left(\left(-\frac{dS(x)^{(0)}}{dx} \frac{dq}{ds}(S(x)^{(0)}) + \frac{dp}{ds}(S(x)^{(0)}) \right) S(x)^{(k)} \right. \\
&\quad - \sum_{i=0}^{N_q} q_i \sum_{\substack{0 \leq j_0, j_1, \dots, j_i < k \\ j_0 + \dots + j_i = k}} \frac{dS(x)^{(j_0)}}{dx} S(x)^{(j_1)} \dots S(x)^{(j_i)} \\
&\quad \left. + \sum_{i=0}^{N_p} p_i \sum_{\substack{0 \leq j_1, \dots, j_i < k \\ j_1 + \dots + j_i = k}} S(x)^{(j_1)} \dots S(x)^{(j_i)} \right). \tag{4.4}
\end{aligned}$$

Now we show that $S(x)^{(k)}$ is an element of $(\text{End}_{\mathbb{C}} M)[[x]]$ by induction on k . The case $k = 0$ follows from $\text{ld}(S(x)^{(0)}) = 1$. For $k > 0$, suppose that $\text{ld}(S(x)^{(k)}) < 0$. This implies $\text{ld}(dS(x)^{(k)}/dx) = \text{ld}(S(x)^{(k)}) - 1$. Since $q_0 \neq 0$ and $S(x)^{(0)}$ is an element of $x\mathbb{C}[[x]]$, $q(S(x)^{(0)})^{-1}$ is an element of $\mathbb{C}[[x]]$. Thus, the lowest degree of the right-hand side of (4.4) is greater than or equal to $\text{ld}(S(x)^{(k)})$ by the induction assumption. This contradicts $\text{ld}(dS(x)^{(k)}/dx) = \text{ld}(S(x)^{(k)}) - 1$. We conclude that $S(x)$ is an element of $(\text{End}_{\mathbb{C}} M)[[x]]$.

We shall show that $S(x)$ is uniquely determined by D and $S_{(0)}$. For all positive integer m , we expand $(dS(x)/dx)q(S(x))$ and $p(S(x))$ modulo

$x^m \mathbb{C}[[x]]$:

$$\begin{aligned}
\frac{dS(x)}{dx} q(S(x)) &\equiv \left(\sum_{j=1}^m j S_{(j)} x^{j-1} \right) \sum_{i=0}^{N_q} q_i \left(\sum_{j=0}^{m-1} S_{(j)} x^j \right)^i \\
&\equiv \sum_{k=0}^{m-1} \sum_{i=0}^{N_q} q_i \sum_{j_0=1}^m \sum_{\substack{0 \leq j_1, \dots, j_i \leq m-1 \\ j_0 + j_1 + \dots + j_i = k+1}} j_0 S_{(j_0)} S_{(j_1)} \cdots S_{(j_i)} x^k \\
&= m S_{(m)} q(S_{(0)}) x^{m-1} \\
&\quad + \sum_{k=0}^{m-1} \sum_{i=0}^{N_q} q_i \sum_{j_0=1}^{m-1} \sum_{\substack{0 \leq j_1, \dots, j_i \leq m-1 \\ j_0 + j_1 + \dots + j_i = k+1}} j_0 S_{(j_0)} S_{(j_1)} \cdots S_{(j_i)} x^k \pmod{x^m \mathbb{C}[[x]]}
\end{aligned} \tag{4.5}$$

and

$$p(S(x)) \equiv \sum_{k=0}^{m-1} \sum_{i=0}^{N_p} p_i \sum_{\substack{0 \leq j_1, \dots, j_i \leq m-1 \\ j_1 + \dots + j_i = k}} S_{(j_1)} \cdots S_{(j_i)} x^k \pmod{x^m \mathbb{C}[[x]]}. \tag{4.6}$$

Comparing the coefficients of x^{m-1} in (4.5) and (4.6), it follows from (4.2) that

$$\begin{aligned}
&m S_{(m)} q(S_{(0)}) \\
&= - \sum_{i=0}^{N_q} q_i \sum_{j_0=1}^{m-1} \sum_{\substack{0 \leq j_1, \dots, j_i \leq m-1 \\ j_0 + j_1 + \dots + j_i = m}} j_0 S_{(j_0)} S_{(j_1)} \cdots S_{(j_i)} \\
&\quad + \sum_{i=0}^{N_p} p_i \sum_{\substack{0 \leq j_1, \dots, j_i \leq m-1 \\ j_1 + \dots + j_i = m-1}} S_{(j_1)} \cdots S_{(j_i)}.
\end{aligned} \tag{4.7}$$

Since $q_0 \neq 0$ and $S_{(0)}$ is nilpotent, $q(S_{(0)})$ is an invertible element of $\text{End}_{\mathbb{C}} M$ and $q(S_{(0)})^{-1}$ is a polynomial in $S_{(0)}$. Thus, it follows by induction on m that every $S_{(m)}$ is a polynomial in $S_{(0)}$ and is uniquely determined by D and $S_{(0)}$. We conclude that $S(x)$ is uniquely determined by D and $S_{(0)}$.

Since every $S_{(m)}, m \in \mathbb{Z}$ is a polynomial in $S_{(0)}$ and M is an indecomposable $\mathcal{A}_M(\mathbb{C}(s))$ -module, the nilpotent element $S_{(0)}$ conjugates to J_n under the identification of $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$. Therefore, Lemma 5 implies that $S(x)$ and hence (M, Y_M) with $\dim_{\mathbb{C}} M = n$ is uniquely determined up to isomorphism under the conditions in (1).

Conversely, suppose that $p(0)q(0) \neq 0$ and $\alpha = p(0)/q(0)$. We shall construct $S(x) \in (\text{End}_{\mathbb{C}} M)[[x]]$ which satisfies the conditions in (1). We identify $\text{End}_{\mathbb{C}} M$ with $\text{Mat}_n(\mathbb{C})$ by fixing a basis of M and we use Lemma 5. Set $S_{(0)} = J_n$. By (4.7) we can inductively define $S_{(m)}$ for $m = 1, 2, \dots$. Reversing the argument used to get (4.7) above, it is easy to see that the obtained upper triangular matrix $S(x) = \sum_{m=0}^{\infty} S_{(m)}x^m \in (\text{Mat}_n(\mathbb{C}))[[x]]$ satisfies (4.2). Taking the semisimple part of (4.7), we obtain the formula by replacing $S_{(j)}$ with $S_{(j)}^{[0]}$ for all $j \in \mathbb{Z}$ in (4.7). By this formula for $m = 1$, we have $S_{(1)}^{[0]} = p_0/q_0 = \alpha$. By this formula again, an inductive argument on m shows $S(x)^{[0]} = \sum_{m=0}^{\infty} S_{(m)}^{[0]}x^m$ is a non-constant element of $\mathbb{C}[[x]]E_n$. This implies that $S(x) - \beta$ is an invertible element of $(\text{Mat}_n(\mathbb{C}))((x))$ for each $\beta \in \mathbb{C}$ and that $\text{ld}(S(x)^{[0]}) = 1$ and $\text{lc}(S(x)^{[0]}) = \alpha$ since $S_{(0)}^{[0]} = 0$. Thus, $q(S(x))$ is an invertible element of $(\text{Mat}_n(\mathbb{C}))((x))$ and hence (4.1) holds. Since all coefficients of $S(x)$ are polynomials in $S_{(0)} = J_n$, $S_{(i)}S_{(j)} = S_{(j)}S_{(i)}$ for all $i, j \in \mathbb{Z}$. By Lemma 5, we have obtained an n -dimensional vertex algebra $(\mathbb{C}(s), D)$ -module M with $\text{ld}(S(x)^{(0)}) = 1$ and with $\text{lc}(S(x)^{(0)}) = \alpha$. This completes the proof of (1).

(2) Next, suppose that $L = 0$. Setting $\tilde{s} = s - \alpha$, we have

$$D = \frac{p(\tilde{s} + \alpha)}{q(\tilde{s} + \alpha)} \frac{d}{d\tilde{s}}.$$

Thus, this case reduces to the case of $L > 0$. It follows by (1) that $S_{(1)}^{[0]} = p(\alpha)/q(\alpha)$.

(3) Finally, suppose that L is a negative integer. Setting $\tilde{s} = 1/s$, we have $Y_M(\tilde{s}, x)^{[0]} = 1/S(x)^{[0]} \in x\mathbb{C}((x))$ and

$$D = \frac{p(1/\tilde{s})}{q(1/\tilde{s})} (-\tilde{s}^2) \frac{d}{d\tilde{s}}.$$

It follows by $S(x)^{[0]} \neq 0$ that $S(x)$ is invertible in $(\text{End}_{\mathbb{C}} M)((x))$ and that $(S(x)^{-1})^{[0]} = (S(x)^{[0]})^{-1}$. Since $S(x)^{-1}$ is a polynomial in $S(x)$, all coefficients in $S(x)^{-1}$ are commutative. Thus, this case also reduces to the case of $L > 0$. $\square$

5 Examples

Throughout this section, D is a non-zero derivation of $\mathbb{C}(s)$ and $f(s) = \sum_{j=0}^m f_j s^j \in \mathbb{C}[s]$ is a square-free polynomial of degree m at least 3. Set $K = \mathbb{C}(s)[t]/(t^2 - f(s))$, which is a quadratic extension of $\mathbb{C}(s)$. Let σ be the generator of the Galois group of K over $\mathbb{C}(s)$ mapping t to $-t$. Since K is a finite Galois extension of $\mathbb{C}(s)$, D can be uniquely extends to a derivation of K , which is also denoted by D .

Let (M, Y_M) be a finite-dimensional indecomposable $(\mathbb{C}(s), D)$ -module. In this section, we shall investigate untwisted or σ -twisted vertex algebra (K, D) -module structures over (M, Y_M) . We denote $Y_M(s, x)$ by $S(x)$ and its semisimple part by $S(x)^{[0]} = \sum_{i=L}^{\infty} S_{(i)}^{[0]} x^i$ with $S_{(L)}^{[0]} \neq 0$ as in Section 4. Theorem 2 says that $L = \text{ld}(S(x)^{[0]}) = 1, 0$, or -1 .

Proposition 6. *Let M be a finite-dimensional indecomposable vertex algebra $(\mathbb{C}(s), D)$ -module and let $(M, \tilde{Y}_M)$ be a g -twisted vertex algebra (K, D) -module structure over (M, Y_M) in Theorem 1, where $g = 1$ or σ . Then*

- (1) *In the case of $\text{ld}(S(x)^{[0]}) = 1$, $g = \sigma$ if and only if $f_0 = 0$.*
- (2) *In the case of $\text{ld}(S(x)^{[0]}) = 0$, $g = \sigma$ if and only if $f(S_{(0)}^{[0]}) = 0$.*
- (3) *In the case of $\text{ld}(S(x)^{[0]}) = -1$, $g = \sigma$ if and only if $\deg f$ is odd.*

Proof. We denote by $\sqrt{f(S(x)^{[0]})}$ a square root of $f(S(x)^{[0]})$ in $\Omega = \cup_{i=1}^{\infty} \mathbb{C}((x^{1/i}))$. We denote by ψ the $\mathbb{C}$ -algebra homomorphism $\psi[\mathbb{C}(s), (M, Y_M)]: \mathbb{C}(s) \rightarrow \mathbb{C}((x))$ defined just after (3.1) and by Q the image of ψ . Let $\hat{K}$ denote the splitting field for $Z^2 - \psi(f(s)) = Z^2 - f(S(x)^{[0]}) \in Q[Z]$ in Ω as in the proof of Theorem 1. We have $\hat{K} \cap \mathbb{C}((x)) = \hat{K}$ if and only if $\sqrt{f(S(x)^{[0]})} \in \mathbb{C}((x))$. By the argument just after (3.3), g is a generator of $\text{Gal}(\hat{K}/\hat{K} \cap \mathbb{C}((x)))$. Thus, $g = 1$ if and only if $\sqrt{f(S(x)^{[0]})} \in \mathbb{C}((x))$. We use the following

expansion of $f(S(x)^{[0]})$:

$$\begin{aligned}
 f(S(x)^{[0]}) &= \sum_{j=0}^m f_j \left(\sum_{i=L}^{\infty} S_{(i)}^{[0]} x^i \right)^j \\
 &= \begin{cases} f_0 + f_1 S_{(1)}^{[0]} x + \cdots, & \text{if } L = 1, \\ f(S_{(0)}^{[0]}) + S_{(1)}^{[0]} f'(S_{(0)}^{[0]}) x + \cdots, & \text{if } L = 0, \\ f_m (S_{(-1)}^{[0]})^m x^{-m} + \cdots, & \text{if } L = -1. \end{cases} \quad (5.1)
 \end{aligned}$$

- (1) In this case if $f_0 \neq 0$, then $\sqrt{f(S(x)^{[0]})} \in \mathbb{C}((x))$. If $f_0 = 0$, then $f_1 \neq 0$ since f is square-free. Thus, $\sqrt{f(S(x)^{[0]})} \notin \mathbb{C}((x))$.
- (2) In this case $S_{(1)}^{[0]} \neq 0$ by Theorem 2 (2). If $f(S_{(0)}^{[0]}) \neq 0$, then $\sqrt{f(S(x)^{[0]})} \in \mathbb{C}((x))$. If $f(S_{(0)}^{[0]}) = 0$, then $\sqrt{f(S(x)^{[0]})} \notin \mathbb{C}((x))$ since f is square-free and $S_{(1)}^{[0]} \neq 0$.
- (3) In this case, the assertion follows easily from (5.1).

□

Acknowledgements

The author thanks the referee for pointing out a mistake in the proof of the earlier version of Theorem 1.

References

- [1] I. Assem, D. Simson, and A. Skowroński, *Elements of the representation theory of associative algebras. Vol. 1*, London Mathematical Society Student Texts, **65**, Cambridge University Press, 2006.
- [2] R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Nat. Acad. Sci. U.S.A.* **83** (1986), 3068–3071.
- [3] C. Dong, H.S. Li and G. Mason, Twisted representations of vertex operator algebras, *Math. Ann.* **310** (1998), 571–600.

- [4] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde, The operator algebra of orbifold models, *Comm. Math. Phys.* **123** (1989), 485–526.
- [5] D. Eisenbud, *Commutative algebra, with a view toward algebraic geometry*, Graduate Texts in Mathematics, **150**, Springer–Verlag, 1995.
- [6] V. G. Kac, *Vertex Algebras for Beginners*, University Lecture Series, **10**, Amer. Math. Soc., 1997.
- [7] J. Lepowsky and H.S. Li, *Introduction to Vertex Operator Algebras and their Representations*, Progress in Mathematics, **227**, Birkhauser Boston, Inc., Boston, MA, 2004.
- [8] R. S. Pierce, *Associative algebras*, Graduate Texts in Mathematics, **88**, Springer–Verlag, 1982.
- [9] K. Tanabe, Finite-dimensional modules for the polynomial ring in one variable as a vertex algebra, *J. Algebra*, **320** (2008), 1261–1274.